



Sussex **MSK** Partnership
Central




Upper Limb Joint Injections in primary care

Workshop – April 2018
Sussex MSK Partnership

+ Tutors

- Giles Hazan – GPwSI MSK, Primary Care Lead SMSK
- Jim Rehill – GP & Primary Care Lead SMSK
- Helen Patten – Advanced Practitioner
- Matt Prout - Advanced Practitioner



PROGRAMME



Time

13.00-13.30 REGISTRATION & LUNCH

13.30-14.40 INTRODUCTION – SAFETY & CONSENT

- SHOULDER INJECTIONS - INDICATIONS & TECHNIQUE
- HAND & WRIST INJECTIONS - INDICATIONS & TECHNIQUE

COFFEE

15.00-16.50

- SHOULDER EXAMINATION & INJECTION PRACTICAL
- HAND/WRIST EXAMINATION & INJECTION PRACTICAL

16.50-17.00 FEEDBACK/CERTIFICATES

+ Contraindications

- Previous Allergy to injection
- Fracture/unstable joint
- Injection into a prosthetic joint
- Uncontrolled coagulopathy
- Ulceration/rash/Infection on skin at injection site
- Injecting into an infected joint or systemic infection
- Injection in the first 16 weeks pregnancy
- Active Psychosis



+ Risks/Complications

- Lipodystrophy
- Loss of skin pigment
- Infection – rare 1:3000-1:50,000 down to 1:2000 in immunosuppressed patients
- Post injection flare
- Facial Flushing (24-72hrs, women>men)
- Altered diabetic control
- Menstrual irregularities
- Fainting



+ Anticoagulation

- WARFARIN: The INR is suggested to be within the therapeutic range and below 4.5 (some suggest below 3)
- NOACs: Newer anticoagulants such as Rivaroxaban and Dabigatran do not need to be stopped but it's worth avoiding the time of peak drug activity e.g. for Rivaroxaban this is 2-4 hrs after the last dose.



+ Consent & Follow Up

- Whilst it may not be medico-legally necessary in England to have a signed consent form it is advised that this should be seen as an adjunct to the clear and sufficient documentation of the discussion of the procedure (including justification and risks) in the notes.
- There is formal guidance available from the GMC and your medical defence organisation may have further information to support you with this.
- It would be of value to audit your injection outcomes to aid future case selection & Appraisal.
- http://www.gmc-uk.org/guidance/ethical_guidance/consent_guidance_index.asp



+ Top 10 tips

1. Have a clear **clinical diagnosis** or diagnostic question in mind before injecting.

2. Tell patients about common **side effects** and also rare but serious side-effects (see leaflet in appendices)

3. Advise patients of the likely **time frame** for improvement and the possibility of short term increase in pain.

4. **Position** yourself and the patient so that you are both comfortable during the procedure

5. Use a good '**no touch**' **technique** for the procedure

6. Have **medication & equipment** available to manage the rare cases of anaphylaxis

7. Have another person available in the clinic in case **help** is needed.

8. Ask the patient to **wait** for about 20-30 minutes before leaving the Clinic

9. **Document** well in your notes - dose, medications, result and any complications (for consent form-see appendices)

10. Consider using a follow up **feedback** form and audit outcomes to guide your case selection.

